

Patient Pocket Guide

A guide to identify signs and symptoms of potential side effects while on immunotherapy, including immune-mediated side effects

1

Detach the wallet card and keep on your person at all times

- Fill out the treating oncologist's contact information

2

Read the information inside and use the Signs and Symptoms Checklist to recognize what to report immediately

- Call your treating oncologist right away if you experience anything unusual while on OPDIVO®, YERVOY®, or Opdualag™
- Contact your oncologist immediately if you experience any symptoms listed on this Signs and Symptoms Checklist, or they get worse. **Getting medical treatment right away may keep the problem from becoming more serious**

3

Show this wallet card during any visits outside of healthcare network

- Tell the urgent care/emergency room staff that you are being treated with OPDIVO, YERVOY, or Opdualag, and show them the important information for Doctors'/Nurses' section of this wallet card

Please see Important Safety Information for OPDIVO, YERVOY, or Opdualag on pages 9–14, a full listing of indications on pages 5–8, and the U.S. Full Prescribing Information and Medication Guide for OPDIVO, YERVOY, and Opdualag.

Please check the box to indicate which immunotherapy the patient is receiving.



OPDIVO
(nivolumab)
INJECTION FOR INTRAVENOUS USE 40 mg/mL



OPDIVO + **YERVOY**
(nivolumab) + (ipilimumab)
INJECTION FOR INTRAVENOUS USE 40 mg/mL INJECTION FOR INTRAVENOUS USE 5 mg/mL



YERVOY
(ipilimumab)
INJECTION FOR INTRAVENOUS INFUSION



Opdualag
(nivolumab and relatlimab-rmbw)
INJECTION FOR INTRAVENOUS USE | 480 mg/160 mg

Detach the wallet card and keep on your person at all times

- Fill out name of immunotherapy drug(s) prescribed by oncologist
- Fill out patient name
- Fill out name and phone number of oncologist
- Keep card in patient's wallet

Be sure to tell all healthcare providers you visit that you are being treated with immunotherapy and SHOW THEM THIS CARD.

Show this wallet card during any urgent care/emergency room visit.

Please call my oncologist.
I have received an immunotherapy called:

Drug(s) Name:

Patient Name:

Oncologist Name:

Oncologist Phone Number:

Present this card **every time** you visit a healthcare provider.

© 2022 Bristol-Myers Squibb Company.

7356-US-2200199 04/22



Please see Important Safety Information for OPDIVO®, YERVOY®, and Opdualag™ on pages 9–14, and the U.S. Full Prescribing Information and Medication Guide for OPDIVO, YERVOY, and Opdualag.

Keep this guide handy for your reference. This guide includes a helpful checklist of Signs and Symptoms for you to closely monitor. Call your Oncology Care Team if you have any of these signs or symptoms, or if they get worse.

These are not all of the possible side effects of OPDIVO® (nivolumab), YERVOY® (ipilimumab) or Opdualag™ (nivolumab and relatlimab-rmbw). This is why you need to report any discomfort or changes in the way you are feeling to your Oncology Care Team.



Heart

- new or worse chest pain
- irregular heartbeat or feel like your heart is racing
- shortness of breath
- tiredness
- swelling in your ankles



Liver

- yellowing of your skin or the whites of your eyes
- severe nausea or vomiting
- pain on the right side of your stomach area (abdomen)
- dark urine (tea colored)
- bleeding or bruising more easily than normal



Intestines

- diarrhea (loose stools) or more frequent bowel movements than usual
- stools that are black, tarry, sticky, or have blood or mucus
- severe stomach-area (abdominal) pain or tenderness




Other Organs

Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with OPDIVO, YERVOY, or Opdualag. Call or see your healthcare provider right away for any new or worsening signs or symptoms, which may include:

- Chest pain, irregular heartbeat, shortness of breath, or swelling of ankles
- Confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs
- Double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight
- Persistent or severe muscle pain or weakness, muscle cramps
- Low red blood cells, bruising

YERVOY may be associated with eye problems with signs and symptoms that include blurry vision, double vision or other vision problems and eye pain or redness.



Hormone Glands

- headaches that will not go away or unusual headaches
- eye sensitivity to light
- eye problems
- rapid heartbeat
- increased sweating
- extreme tiredness
- weight gain or weight loss
- feeling more hungry or thirsty than usual
- urinating more often than usual
- hair loss
- feeling cold
- constipation
- your voice gets deeper
- dizziness or fainting
- changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness



Lungs

- new or worsening cough
- chest pain
- shortness of breath



Kidneys

- decrease in your amount of urine
- blood in your urine
- swelling of your ankles
- loss of appetite



Skin

- rash
- itching
- skin blistering or peeling
- painful sores or ulcers in mouth or nose, throat, or genital area



Severe Infusion-Related Reactions

- chills or shaking
- itching or rash
- flushing
- shortness of breath or wheezing
- dizziness
- feel like passing out
- fever
- back or neck pain

Do not treat symptoms yourself

Symptoms may occur anytime during treatment or even after your treatment has ended

Do not feel embarrassed or that you are bothering your oncologist

Complications, including graft-versus-host disease (GVHD), of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic). These complications can be severe and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with OPDIVO, YERVOY, or Opdualag. Your healthcare provider will monitor you for these complications.

Please see Important Safety Information for OPDIVO, YERVOY, and Opdualag on pages 9–14, and the U.S. Full Prescribing Information and Medication Guide for OPDIVO, YERVOY, and Opdualag.

Important Safety Information

OPDIVO® (nivolumab), YERVOY® (ipilimumab), and Opdualag™ (nivolumab and relatlimab-rmbw) are medicines that may treat certain cancers by working with your immune system. OPDIVO, YERVOY, and Opdualag can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. These problems may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time. Some of these problems may happen more often when OPDIVO is used in combination with YERVOY or another therapy.

YERVOY can cause serious side effects in many parts of your body which can lead to death. These problems may happen anytime during treatment with YERVOY or after you have completed treatment

OPDIVO® (nivolumab) and OPDIVO + YERVOY® (ipilimumab) Indications

-  OPDIVO® (nivolumab) is a prescription medicine used to treat adults with a type of skin cancer called melanoma that has spread or cannot be removed by surgery (advanced melanoma).
-  OPDIVO® (nivolumab) is a prescription medicine used in combination with YERVOY® (ipilimumab) to treat adults with a type of skin cancer called melanoma that has spread or cannot be removed by surgery (advanced melanoma).
-  OPDIVO® (nivolumab) is a prescription medicine used in combination with chemotherapy that contains platinum and another chemotherapy medicine, before you have surgery, for adults with early-stage lung cancer (called non-small cell lung cancer).
-  OPDIVO® (nivolumab) is a prescription medicine used in combination with YERVOY® (ipilimumab) as a first treatment for adults with a type of advanced stage lung cancer (called non-small cell lung cancer) when your lung cancer has spread to other parts of your body (metastatic) **and** your tumors are positive for PD-L1, but do not have an abnormal EGFR or ALK gene.
-  OPDIVO® (nivolumab) is a prescription medicine used in combination with YERVOY® (ipilimumab) and 2 cycles of chemotherapy that contains platinum and another chemotherapy medicine, as a first treatment for adults with a type of advanced stage lung cancer (called non-small cell lung cancer) when your lung cancer has spread or grown, or comes back, **and** your tumor does not have an abnormal EGFR or ALK gene.
-  OPDIVO® (nivolumab) is a prescription medicine used to treat adults with a type of advanced stage lung cancer (called non-small cell lung cancer) that has spread or grown **and** you have tried chemotherapy that contains platinum, and it did not work or is no longer working. If your tumor has an abnormal EGFR or ALK gene, you should have also tried an FDA-approved therapy for tumors with these abnormal genes, and it did not work or is no longer working.

-  OPDIVO® (nivolumab) is a prescription medicine used in combination with YERVOY® (ipilimumab) as a first treatment for adults with a type of cancer that affects the lining of the lungs and chest wall called malignant pleural mesothelioma which cannot be removed by surgery.
-  OPDIVO® (nivolumab) is a prescription medicine used in combination with YERVOY® (ipilimumab) to treat adults with kidney cancer in certain people when your cancer has spread (advanced renal cell carcinoma) and you have not already had treatment for your advanced RCC.
-  OPDIVO® (nivolumab) is a prescription medicine used in combination with cabozantinib to treat adults with kidney cancer when your cancer has spread (advanced renal cell carcinoma) and you have not already had treatment for your advanced RCC. Please read the Patient Information that comes with cabozantinib.
-  OPDIVO® (nivolumab) is a prescription medicine used to treat adults with kidney cancer (renal cell carcinoma) when your cancer has spread or grown after treatment with other cancer medications.
-  OPDIVO® (nivolumab) is a prescription medicine used to treat adults with a type of blood cancer called classical Hodgkin lymphoma if your cancer has come back or spread after a type of stem cell transplant that uses your own stem cells (autologous), **and** you used the medicine brentuximab vedotin before or after your stem cell transplant, **or** if you received at least 3 kinds of treatment including an autologous stem cell transplant. OPDIVO was approved based on response rate. There is ongoing evaluation of clinical benefit of OPDIVO for this use.
-  OPDIVO® (nivolumab) is a prescription medicine used to treat adults with head and neck cancer (squamous cell carcinoma) that has come back or spread **and** you have tried chemotherapy that contains platinum and it did not work or is no longer working.
-  OPDIVO® (nivolumab) is a prescription medicine used to treat adults with cancer of the lining of the urinary tract (including the bladder, urethra, ureters, or renal pelvis) [urothelial carcinoma] when it has spread or grown (locally advanced or metastatic) and you have tried chemotherapy that contains platinum, and it did not work or is no longer working or your cancer worsened within 12 months of treatment with chemotherapy that contains platinum, either before or after surgery to remove your cancer.
-  OPDIVO® (nivolumab) is a prescription medicine used to treat adults with cancer of the lining of the urinary tract (including the bladder, ureters, or renal pelvis) [urothelial carcinoma] to help prevent cancer of the urinary tract from coming back after it was removed by surgery.

Please see Important Safety Information for OPDIVO, YERVOY, and Opdualag on pages 9–14, and the U.S. Full Prescribing Information and Medication Guide for [OPDIVO](#), [YERVOY](#), and [Opdualag](#).

OPDIVO® (nivolumab) and OPDIVO + YERVOY® (ipilimumab) Indications (cont'd)

-  OPDIVO® (nivolumab) is a prescription medicine used to treat adults and children 12 years of age and older with a type of colon or rectal cancer (colorectal cancer) that has spread to other parts of the body (metastatic), is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR), **and** you have tried treatment with a fluoropyrimidine, oxaliplatin, and irinotecan, and it did not work or is no longer working. OPDIVO was approved based on response rate and how long patients' responses lasted. There is ongoing evaluation of clinical benefit of OPDIVO for this use.
-  OPDIVO® (nivolumab) is a prescription medicine used in combination with YERVOY® (ipilimumab) to treat adults and children 12 years of age and older, with a type of colon or rectal cancer (colorectal cancer) that has spread to other parts of the body (metastatic), is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR), **and** you have tried treatment with a fluoropyrimidine, oxaliplatin, and irinotecan, and it did not work or is no longer working. OPDIVO in combination with YERVOY was approved based on response rate and how long patients' responses lasted. There is ongoing evaluation of clinical benefit of OPDIVO in combination with YERVOY for this use.
-  OPDIVO® (nivolumab) is a prescription medicine used in combination with YERVOY® (ipilimumab) to treat adults with liver cancer (hepatocellular carcinoma) if you have previously received treatment with sorafenib. OPDIVO in combination with YERVOY was approved based on response rate and how long patients' responses lasted. There is ongoing evaluation of clinical benefit of OPDIVO in combination with YERVOY for this use.
-  OPDIVO® (nivolumab) is a prescription medicine used to treat adults with a type of skin cancer called melanoma to help prevent melanoma from coming back after it and lymph nodes that contain cancer have been removed by surgery.
-  OPDIVO® (nivolumab) is a prescription medicine used to treat adults with cancer of the tube that connects your throat to your stomach (esophageal cancer) if your esophageal cancer is a type called squamous cell carcinoma, **and** cannot be removed with surgery, **and** has come back or spread to other parts of the body after you have received chemotherapy that contains fluoropyrimidine and platinum.
-  OPDIVO® (nivolumab) is a prescription medicine that may be used in adults to help prevent cancer of the tube that connects your throat to your stomach (esophageal cancer) or cancer where the esophagus joins the stomach (gastroesophageal junction cancer) from coming back after it has been treated with chemoradiation followed by surgery to completely remove the cancer, **but** some cancer cells were still present in the removed tumor or lymph nodes.

-  OPDIVO® (nivolumab) is a prescription medicine used in combination with chemotherapy that contains fluoropyrimidine and platinum to treat adults with cancer of the stomach (gastric), junction between the stomach and esophagus (gastroesophageal junction), and esophagus that is a type called adenocarcinoma and cannot be removed with surgery **or** has spread to other parts of the body.

It is not known if OPDIVO is safe and effective in children younger than 12 years of age with MSI-H or dMMR metastatic colorectal cancer, **or** in children younger than 18 years of age for the treatment of any other cancers.

YERVOY Indications

-  YERVOY® (ipilimumab) is a prescription medicine used in adults and children 12 years of age and older to treat melanoma (a kind of skin cancer) that has spread (metastatic) or cannot be removed by surgery (unresectable).

It is not known if YERVOY is safe and effective in children less than 12 years of age.
-  YERVOY® (ipilimumab) is a prescription medicine used in adults to help prevent melanoma (a kind of skin cancer) from coming back after it and lymph nodes that contain cancer have been removed by surgery.

It is not known if YERVOY is safe and effective in children.

Opdualag Indication

-  Opdualag™ (nivolumab and relatlimab-rmbw) is a prescription medicine used to treat adults and children 12 years of age or older with a type of skin cancer called melanoma that has spread or cannot be removed by surgery (advanced melanoma).

It is not known if Opdualag is safe and effective when used in children younger than 12 years of age or older who weigh less than 88 pounds (40 kg), or in children younger than 12 years of age.

Please see Important Safety Information for OPDIVO, YERVOY, and Opdualag on pages 9–14, and the U.S. Full Prescribing Information and Medication Guide for [OPDIVO](#), [YERVOY](#), and [Opdualag](#).

Important Safety Information for OPDIVO® (nivolumab), YERVOY® (ipilimumab), and OPDIVO + YERVOY

What is the most important information I should know about OPDIVO and YERVOY?

OPDIVO and YERVOY are medicines that may treat certain cancers by working with your immune system. OPDIVO and YERVOY can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. These problems may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time. Some of these problems may happen more often when OPDIVO is used in combination with YERVOY or another therapy

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

- **Lung problems:** new or worsening cough; shortness of breath; chest pain
- **Intestinal problems:** diarrhea (loose stools) or more frequent bowel movements than usual; stools that are black, tarry, sticky, or have blood or mucus; severe stomach-area (abdominal) pain or tenderness
- **Liver problems:** yellowing of your skin or the whites of your eyes; severe nausea or vomiting; pain on the right side of your stomach area (abdomen); dark urine (tea colored); bleeding or bruising more easily than normal
- **Hormone gland problems:** headaches that will not go away or unusual headaches; eye sensitivity to light; eye problems; rapid heart beat; increased sweating;

extreme tiredness; weight gain or weight loss; feeling more hungry or thirsty than usual; urinating more often than usual; hair loss; feeling cold; constipation; your voice gets deeper; dizziness or fainting; changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness

- **Kidney problems:** decrease in your amount of urine; blood in your urine; swelling in your ankles; loss of appetite
- **Skin problems:** rash; itching; skin blistering or peeling; painful sores or ulcers in the mouth or nose, throat, or genital area
- **Eye problems:** blurry vision, double vision, or other vision problems; eye pain or redness

Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with OPDIVO and YERVOY. Call or see your healthcare provider right away for any new or worsening signs or symptoms, which may include:

- Chest pain; irregular heartbeat; shortness of breath; swelling of ankles
- Confusion; sleepiness; memory problems; changes in mood or behavior; stiff neck; balance problems; tingling or numbness of the arms or legs
- Double vision; blurry vision; sensitivity to light; eye pain; changes in eyesight
- Persistent or severe muscle pain or weakness; muscle cramps
- Low red blood cells; bruising

Getting medical help right away may help keep these problems from becoming more serious. Your healthcare team will check you for these problems during treatment and may treat you with corticosteroid or hormone replacement medicines. Your healthcare team may also need to delay or completely stop your treatment if you have severe side effects.

Possible side effects of OPDIVO and YERVOY

OPDIVO and YERVOY can cause serious side effects, including:

- **See “What is the most important information I should know about OPDIVO and YERVOY?”**
- **Severe infusion reactions.** Tell your healthcare team right away if you get these symptoms during an infusion of OPDIVO or YERVOY: chills or shaking; itching or rash; flushing; shortness of breath or wheezing; dizziness; feel like passing out; fever; back or neck pain
- **Complications, including graft-versus-host disease (GVHD), of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic).** These complications can be severe and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with OPDIVO or YERVOY. Your healthcare provider will monitor you for these complications.

The most common side effects of OPDIVO, when used alone, include: feeling tired; rash; pain in muscles, bones, and joints; itchy skin; diarrhea; nausea; weakness; cough; vomiting; shortness of breath; constipation; decreased appetite;

back pain; upper respiratory tract infection; fever; headache; stomach-area (abdominal) pain; and urinary tract infection

The most common side effects of OPDIVO, when used in combination with chemotherapy, include: nausea, constipation, feeling tired, decreased appetite, and rash.

The most common side effects of OPDIVO, when used in combination with YERVOY, include: feeling tired; diarrhea; rash; itching; nausea; pain in muscles, bones, and joints; fever; cough; decreased appetite; vomiting; stomach-area (abdominal) pain; shortness of breath; upper respiratory tract infection; headache; low thyroid hormone levels (hypothyroidism); constipation; decreased weight; and dizziness.

The most common side effects of OPDIVO, when used in combination with YERVOY and chemotherapy, include: feeling tired; pain in muscles, bones, and joints; nausea; diarrhea; rash; decreased appetite; constipation; and itching.

The most common side effects of OPDIVO, when used in combination with cabozantinib, include: diarrhea; feeling tired or weak; liver problems; rash, redness, pain, swelling or blisters on the palms of your hands or soles of your feet; mouth sores; rash; high blood pressure; low thyroid hormone levels; pain in muscles, bones, and joints; decreased appetite; nausea; change in the sense of taste; stomach-area (abdominal) pain; cough; upper respiratory tract infection.

Please see Important Safety Information for [OPDIVO](#), [YERVOY](#), and [Opdualag](#)™ (nivolumab and relatlimab-rmbw) continued on the following page.

Important Safety Information for OPDIVO® (nivolumab), YERVOY® (ipilimumab), and OPDIVO + YERVOY (continued)

The most common side effects of OPDIVO when used in combination with fluoropyrimidine and platinum-containing chemotherapy include: numbness, pain, tingling, or burning in your hands or feet; nausea; feeling tired; diarrhea; vomiting; decreased appetite; stomach-area (abdominal) pain; constipation; pain in muscles, bones, and joints.

The most common side effects of YERVOY include: feeling tired; diarrhea; nausea; itching; rash; vomiting; headache; weight loss; fever; decreased appetite; difficulty falling or staying asleep

These are not all the possible side effects. For more information, ask your healthcare provider or pharmacist. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.

Before receiving OPDIVO or YERVOY, tell your healthcare provider about all of your medical conditions, including if you:

- have immune system problems such as Crohn's disease, ulcerative colitis, or lupus
- have received an organ transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have received radiation treatment to your chest area in the past and have received other medicines that are like OPDIVO

- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome
- are pregnant or plan to become pregnant. OPDIVO and YERVOY can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if OPDIVO or YERVOY passes into your breast milk. Do not breastfeed during treatment with OPDIVO or YERVOY and for 5 months after the last dose of OPDIVO and 3 months after the last dose of YERVOY.

Females who are able to become pregnant:

Your healthcare provider should do a pregnancy test before you start receiving OPDIVO or YERVOY.

- You should use an effective method of birth control during your treatment and for at least 5 months after the last dose of OPDIVO and for 3 months after the last dose of YERVOY. Talk to your healthcare provider about birth control methods that you can use during this time.
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with OPDIVO or YERVOY. You or your healthcare provider should contact Bristol Myers Squibb at 1-844-593-7869 as soon as you become aware of a pregnancy.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Important Safety Information for Opdualag™ (nivolumab and relatlimab-rmbw)

What is the most important information I should know about Opdualag?

Opdualag is a medicine that may treat a type of skin cancer called melanoma by working with your immune system. Opdualag can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or can lead to death. These problems may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

- **Lung problems:** new or worsening cough; shortness of breath; chest pain
- **Intestinal problems:** diarrhea (loose stools) or more frequent bowel movements than usual; stools that are black, tarry, sticky, or have blood or mucus; severe stomach-area (abdominal) pain or tenderness
- **Liver problems:** yellowing of your skin or the whites of your eyes; severe nausea or vomiting; pain on the right side

of your stomach area (abdomen); dark urine (tea colored); bleeding or bruising more easily than normal

- **Hormone gland problems:** headaches that will not go away or unusual headaches; eye sensitivity to light; eye problems; rapid heartbeat; increased sweating; extreme tiredness; weight gain or weight loss; feeling more hungry or thirsty than usual; urinating more often than usual; hair loss; feeling cold; constipation; your voice gets deeper; dizziness or fainting; changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness
- **Kidney problems:** decrease in your amount of urine; blood in your urine; swelling in your ankles; loss of appetite
- **Skin problems:** rash; itching; skin blistering or peeling; painful sores or ulcers in the mouth or nose, throat, or genital area
- **Heart problems:** new or worse chest pain; irregular heartbeat or feel like your heart is racing; shortness of breath; tiredness; swelling in your ankles

Please see Important Safety Information for [OPDIVO](#), [YERVOY](#), and [Opdualag](#) continued on the following page.

Important Safety Information for Opdualag™ (nivolumab and relatlimab-rmbw) (continued)

Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with Opdualag. Call or see your healthcare provider right away for any new or worsening signs or symptoms, which may include:

- Confusion; sleepiness; memory problems; changes in mood or behavior; stiff neck; balance problems; tingling or numbness of the arms or legs
- Double vision; blurry vision; sensitivity to light; eye pain; changes in eyesight
- Persistent or severe muscle pain or weakness; muscle cramps
- Low red blood cells; bruising

Getting medical help right away may help keep these problems from becoming more serious. Your healthcare team will check you for these problems during treatment with Opdualag. Your healthcare provider may treat you with corticosteroid or hormone replacement medicines. Your healthcare provider may also need to delay or completely stop your treatment with Opdualag if you have severe side effects.

Possible side effects of Opdualag

Opdualag can cause serious side effects, including:

- See “What is the most important information I should know about Opdualag?”

- **Severe infusion reactions.** Tell your healthcare provider or nurse right away if you get these symptoms during an infusion of Opdualag: chills or shaking; itching or rash; flushing; shortness of breath; dizziness; feel like passing out; fever; back or neck pain
- **Complications of stem cell transplant that uses donor stem cells (allogeneic).** These complications can be severe and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with Opdualag. Your healthcare provider will monitor you for signs of complications if you have an allogeneic stem cell transplant.

The most common side effects of Opdualag, include: muscle and bone pain; tiredness; decreased red blood cell and white blood cell counts; increased liver function test results; rash; itching, diarrhea; decreased salt (sodium) in your blood

These are not all the possible side effects of Opdualag. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Before receiving Opdualag, tell your healthcare provider if you:

- have immune system problems such as Crohn’s disease, ulcerative colitis, or lupus
- have received an organ transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have a condition that affects your nervous system, such as myasthenia gravis, or Guillain-Barré syndrome
- are pregnant or plan to become pregnant. Opdualag can harm your unborn baby

Females who are able to become pregnant:

- your healthcare provider should do a pregnancy test before you start receiving Opdualag
- you should use an effective method of birth control during and for at least 5 months after the last dose of Opdualag. Talk to your healthcare provider about birth control methods that you can use during this time
- Tell your healthcare provider right away if you become pregnant during treatment with Opdualag

- are breastfeeding or plan to breastfeed. It is not known if Opdualag passes into your breast milk. Do not breastfeed during treatment with Opdualag and for 5 months after the last dose of Opdualag

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

IMPORTANT Reminders for Patients

- Contact your oncologist immediately if you experience any symptoms listed on this Signs and Symptoms Checklist or if they get worse, or if you experience anything unusual while on OPDIVO® (nivolumab), YERVOY® (ipilimumab), or Opdualag™ (nivolumab and relatlimab-rmbw) therapy. Getting medical treatment right away may keep the problem from becoming more serious
- These are not all of the possible side effects of OPDIVO, YERVOY, or Opdualag. This is why you need to report any discomfort or changes in the way you are feeling to your Oncology Care Team
- Do not treat Signs and Symptoms yourself without talking to your oncologist
- Your oncologist may decide to delay or completely stop OPDIVO, OPDIVO + YERVOY, YERVOY, or Opdualag, or give you other medicines to treat your symptoms

Please see Important Safety Information for OPDIVO, YERVOY, and Opdualag on pages 9–14, and the U.S. Full Prescribing Information and Medication Guide for OPDIVO, YERVOY, and Opdualag.

Live On-Demand Support Services for You and Your Patients

We're here to help you better understand your medication. These programs are for people who are prescribed OPDIVO or Opdualag and their loved ones. Signing up takes less than 5 minutes. Bristol Myers Squibb remains committed to responding to your needs within 24 hours or less.

OPDIVO with You
(nivolumab)
INJECTION FOR INTRAVENOUS USE 10 mg/mL



1-855-OPDIVO-1
(1-855-673-4861)

FOR LIVE SUPPORT AND ASSISTANCE
8:00 AM to 8:00 PM ET, Monday-Friday

Responses provided between
8:00 AM to 8:00 PM ET, Monday-Friday

- Helpful information
- Details about support and advocacy groups
- A care counselor as close as your phone

These support resources and more can be found on
www.opdivo.com

Opdualag with You
(nivolumab and relatimab-mbw)
Injection for intravenous use | 480 mg/160 mg



Helpful Information

Get useful tips, tools, and educational materials that can help you or your loved one learn more about Opdualag and may help support your care.



Support and Advocacy Groups

We can direct you to organizations that may be helpful for the challenges you may face.



Just call **1-855-OPDUALAG** (1-855-673-8252)
8:00 AM to 8:00 PM ET, Monday-Friday

These support resources and more can be found on
www.Opdualag.com

Please see Important Safety Information for OPDIVO, YERVOY, and Opdualag on pages 9–14, and the U.S. Full Prescribing Information and Medication Guide for OPDIVO, YERVOY, and Opdualag.



OPDIVO®, YERVOY®, Opdualag™ and the related logos are trademarks of Bristol-Myers Squibb Company. Other brands listed are the trademarks of their respective owners.
© 2022 Bristol-Myers Squibb Company.
7356-US-2200199 04/22

OPDIVO
(nivolumab)
INJECTION FOR INTRAVENOUS USE 10 mg/mL

OPDIVO + **YERVOY**
(nivolumab) (ipilimumab)

YERVOY
(ipilimumab)
Injection for intravenous infusion

Opdualag
(nivolumab and relatimab-mbw)
Injection for intravenous use | 480 mg/160 mg

Your Signs and Symptoms Checklist

Do you have any signs or symptoms?

1. If you notice any discomfort or changes in the way you are feeling, contact your Oncology Care Team immediately.
 2. Getting medical treatment right away may help keep these problems from becoming more serious. Do not try to treat possible side effects yourself.
 3. Your oncologist may decide to delay or completely stop OPDIVO® (nivolumab), YERVOY® (ipilimumab), or Opdualag™ (nivolumab and relatlimab-rmbw) or give you other medicines to treat your symptoms.
- **These are not all of the possible side effects of OPDIVO, YERVOY, or Opdualag. This is why you need to report any discomfort or changes in the way you are feeling to your Oncology Care Team. Signs or changes to look out for include:**



Liver

- yellowing of your skin or the whites of your eyes
- severe nausea or vomiting
- pain on the right side of your stomach area (abdomen)
- dark urine (tea colored)
- bleeding or bruising more easily than normal



Intestines

- diarrhea (loose stools) or more frequent bowel movements than usual
- stools that are black, tarry, sticky, or have blood or mucus
- severe stomach-area (abdominal) pain or tenderness



Lungs

- new or worsening cough
- chest pain
- shortness of breath



Skin

- rash
- itching
- skin blistering or peeling
- painful sores or ulcers in mouth or nose, throat, or genital area



Kidneys

- decrease in your amount of urine
- blood in your urine
- swelling of your ankles
- loss of appetite



Severe Infusion-Related Reactions

- chills or shaking
- itching or rash
- flushing
- shortness of breath or wheezing
- dizziness
- feel like passing out
- fever
- back or neck pain



Hormone Glands

- headaches that will not go away or unusual headaches
- eye sensitivity to light
- eye problems
- rapid heartbeat
- increased sweating
- extreme tiredness
- weight gain or weight loss
- feeling more hungry or thirsty than usual
- urinating more often than usual
- hair loss
- feeling cold
- constipation
- your voice gets deeper
- dizziness or fainting
- changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness



Heart

- new or worse chest pain
- irregular heartbeat or feel like your heart is racing
- shortness of breath
- tiredness
- swelling in your ankles



Other Organs

- Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with OPDIVO, YERVOY, or Opdualag. Call or see your healthcare provider right away for any new or worsening signs or symptoms, which may include:
- Chest pain, irregular heartbeat, shortness of breath or swelling of ankles
 - Confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs
 - Double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight
 - Persistent or severe muscle pain or weakness, muscle cramps
 - Low red blood cells, bruising

YERVOY may be associated with eye problems with signs and symptoms that include blurry vision, double vision, or other vision problems and eye pain or redness.

Complications, including graft-versus-host disease (GVHD), of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic). These complications can be severe and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with OPDIVO, YERVOY, or Opdualag. Your healthcare provider will monitor you for these complications.

If you experience a severe side effect, you may need to go to a *hospital emergency department*. If this happens, you need to:



- Bring the wallet card and a full list of your medications, including prescription and over-the-counter medicines, vitamins, and herbal supplements
- Tell doctors and nurses in the emergency department that **you are on an immunotherapy treatment** to treat your condition
- Make sure your Oncology Care Team is notified

Please see Important Safety Information for OPDIVO, YERVOY, and Opdualag on pages 9–14, and the U.S. Full Prescribing Information and Medication Guide for OPDIVO, YERVOY, and Opdualag.

Management of immunotherapy side effects may require a different approach than other cancer treatments. Do not wait to treat the patient if urgent care is required.

Call patient's oncologist.

ATTENTION



IMPORTANT Information for Doctors/Nurses

Please call my oncologist.

I have received an immunotherapy called:

Drug(s) Name:

Patient Name:

Oncologist Name:

Oncologist Phone Number:

Present this card **every time** you visit a healthcare provider.